THE EFFECT OF ANTIHISTAMINIC DRUG ON TUBERCULIN REACTION

2

EUNG SOO HAN, M.D.

Seoul, Korea

Ministry of Health,

INTRODUCTION

It is well known that antihistaminic drugs have a marked effect on the manifestations of allergy. Considerable interest has been raised as to the effect that this set of drugs could have on the intradermal tuberculin reaction in man and animals. Many workers have studied the subject recently in this respect, but the results of their investigations have been divergent and have created sharp controversies.

Bouquet (1) and Sarber (2) have made investigations on animals. Bouquet (1) obtained on appreciable inhibitory effect on the tuberculin reaction of guinea pigs by using different kinds of antihistaminics. On the other hand, Sarber (2) obtained marked reduction of the tuberculin reaction both in size and intensity.

This controversy evoked further interest which resulted in the clinical investigations by Henderson and Jodd (3) and Henderson (4). The latter workers gave large doses of Pyribenzamine or similar groups of drugs daily for as long

as six months and claimed that inhibitory effect was obtained. They went one step further to say that it was possible to produce marked clinical improvement in tuberculous pneumonia and exudative lesions in this way. They employed 30 such cases for their study and of these 18 cases showed improvement.

Suppression of the tuberculin reaction, they found, was reversed within 4-6 weeks after the discontinuation of the antihistaminic drugs. Graub and Barrist (5) supported the above results and concluded that the tuberculin testing should not be done when antihistaminic drugs are being administered. The opposite conclusion was arrived at by Cripe, Levine and Aaron (6) and Friedman and Silberman (7). Hunter, Davis and Hyde (8) mentioned that antihistaminic drugs such as pyribenzamine and Benadryl produced no effect whatsoever either on the tuberculin reaction or on the pulmonary process.

In a recent letter to the editors of American Review of Tuberculosis, (9) made the suggestion "one would not expect drugs recommended for use in the

Foot Note; 1. This experimental study was made in February 1951 at the North Carolina Sanatorium, McCain, N.C., U.S.A.

2. Tuberculosis Controller,

Republic of Korea

3. This paper was read at the First Medical Session, Korean Tuberculosis Association, Seoul in November 7, 1953

nylactic type of allergic reaction to have much effect on the delayed, tuberculin type of reaction."

Because of the varied opinions and controversies existing on the subject, re-investigation appeared worth while. The conclusion on this subject should be of great importance because of the extensive use of these drugs (10 a,b.) particularly during winter and spring.

PART I PRELIMINARY EXPERIMENT

A preliminary experiment was conducted to determine the experimental dosage of antihistaminics as well as to observe clinical side effects following administration of various amount of antihistaminics.

METHODS OF INVESTIGATION

Antihistaminic drugs: Pyribenzamine tablet (N, N-Dimethyl-N¹-benzyl-N¹(alpha-pyridyl) ethylene diamine hydrochloride. Ciba Pharmaceutical Products, 50 mgm, each.

Patients; These are all patients in different phases of proved tuberculosis at the North Carolina Sanatorium, McCain, N.C. Tuberculin Test; Old Tuberculin was used throughout the entire study. Old Tuberculin was diluted in physiologic salt solution, 1--10,000. The technique and interpretation as outlined by the National Tuberculosis Association, 1950, were followed. Reaction and diameter of redness and induration were carefully recorded in millimeters. Any decrease of more than 5 millimeters in diameter was considered as inhibition.

Dosages: Large dose group received 400 mgm a day in four divided doses.

Medium dose group received 200 mgm a day in four divided doses.

Small dose group received 50 mgm a day.

Experiments were divided into two major groups as follows:

I. Pre-medicated Group

Antihistaminics given prior to tuberculin test.

a. Long term group: A 7-day period of medication prior to skin test and continued for 48 hours while observing skin test.

1. Large dose group;
2. Medium dose group;
3. Small dose group;

b. Short term group: A 2-day period of medication prior to the skin test and continued for 48 hours while observing skin test.

1. Large dose group;
2. Medium dose group;
3. Small dose group;

II. Non-Pre-Medicated Group

Antihistaminics were not given prior to skin test but given with skin test and continued for 48 hours while skin test was observed.

1. Large dose group;
2. Medium dose group;
3. Small dose group;

These patients had not received any kind of antihistaminic drugs for a period of at least three weeks prior to the investigation. However, individual patients received their routine regimen or drug therapy such as Streptomycin or Para-Aminosalicylic Acid compounds regardless of this observation.

RESULTS

1. Pre-Medicated Group

Table 1 and 2 show the results of long and short term groups respectively.

TABLE 1

Long term group

Name of pati.   Results (48 hours)
& Dosage gro.   Before Med.   After Med.

a. Large dose group

J.N.	5x8	15x18
w.J.	15x18	13x14/40x45
S.H.	12x12	12x16

b. Medium dose group

B.W.	10x10	8x10
T.A.	12x12	32x35
A.G.	11x13	10x10

c. Small dose group

H.J.	12x12	14x15
D.S.	22x25	7x14/70x90
R.S.	12x12	14x15

There were no appreciable inhibitory effects after the medication regardless of the dosage of antihistaminic drugs used in this series.

TABLE II

Short term group

Name of pati.   Results (48 hours)
& Dosage gro.   Before Med.   After Med.

a. Large dose group

G.M.	15x15	20x22
A.W.	12x16	8x10
H.J.	10x10/15x22	14x15

b. Medium dose group

P.O.	12x12/24x38	17x20/25x45
D.J.	12x14	15x15
C.W.	6x6	1010x

c. Small dose group

W.E.	22x24	15x21/25x45
L.G.	6x6	10x12
K.D.	7x7	6x6/12x18

In the large dose group two of three cases showed remarkable reduction in their size of reaction, though one case showed increase in size of reaction. However, there were no appreciable inhibitory effects among the groups which received medium and small doses.

II. Non-pre-Medicated Group

Table 3 shows that no appreciable inhibitory effects regardless of the dosages, except one case in the small dose group M. R.

TABLE 3

Name of pati.   Results (48 hours)
& Dosage gro.   Before Med.   After Med.

a. Large dose group

T.T.	12x13	12x15
D.L.	12x12	14x16
H.F.	13x13/45x50	17x18/45x55

b. Medium dose group

L.S.	8x8	14x15
D.R.	12x17	18x24
E.W.	12x13	17x20

c. Small dose group

C.W.	10x10	13x15
M.R.	15x16/42x60	22x25
D.C.	15x16	18x20

SIDE REACTIONS OF...

...ANTIHISTAMINICS

Table 4 shows the side reactions caused by the antihistaminic drugs with the different doses used. Except one case, throughout the study the antihistaminic used was Pyribenzamine. One case T. T. was treated with Benadryl as he developed severe reactions within one hour after

er the first dose of a 100 mgm. of Pyridenzamine. However, these side reactions mentioned below in the table were mostly limited to the large dose group who received daily 400 mgm., but a few among the group who received medium dose (so-called clinical dosage) of a 200 mgm. daily also showed uncomfortable side reactions to a minor degree. No such side reactions were seen in the small dose group who received daily dosage of 50 mgm.

These side reactions promptly disappeared after the discontinuation of medication with no undesirable symptoms.

TABLE 4

Symp- toms	Daily dosage	Degree of reaction	No. of Cases	Percent
Vertigo	400mgm.	Severe	3	27.3%
		Moderately		
		Severe	2	18.2%
	200mgm.	Mild	3	27.3%
		Moderately		
		Severe	1	11.1%
Rest- less- ness	400mgm.	Mild	3	33.3%
		Severe		
		moderately sever	2	18.2%
	200mgm.	Mild	1	9.0%
		moderately		
		severe	3	11.1%
Head- ache	400mgm.	Severe	3	27.3%
		Mild	1	9.0%
Anor- exia & Nausea	200mgm.	Mild	1	11.1%
	400mgm.	above 100°F	1	9.0%

DISCUSSION

Pyridenzamine in large doses produced undesirable clinical side reactions such as dizziness, headache, restlessness, anorexia and high fever but those symptoms promptly disappeared shortly after the discontinuation of the medication. It was found that 200 mgm. daily

appeared to be adequate to conduct this investigation.

It was noted that Pyridenzamine does not play any significant role in influencing the tuberculin reaction one way or the other. However, a 400mgm. daily dose of Pyridenzamine for a short period of two days prior to the skin test showed marked inhibitory effect. The number of patients investigated in this series are few and the knowledge arrived at through such a limited number of experiments can be contributory only in a small measure to a final conclusion. Therefore, a further investigation using a 200 mgm. of Pyridenzamine daily on a large number of patients is planned as following outlined;

PART II. MAIN EXPERIMENT

Experimental groups were divided into three major groups as outlined in the preliminary study; Pre-Medicated Group of long and short term, Non-Pre-Medicated group and Control Group of non-medicated but tuberculin test was repeated along with the experimental groups.

P.P.D. (Purified Protein Derivatives), Sahrp & Dohme No. 4481, first strength, 0.0002 mgm was used instead of Old Tuberculin and reaction was measured in millimeters and otherwise, the experimental methods are exactly followed the preliminary study. In addition to the above described, cough and sputum were carefully studied to observe the patients clinical course following medication.

RESULTS

1. PRE-MEDICATED GROUP

Table 1 and 2 show the results of

long term and short term groups respectively.

Table 1: Long term group

Name	Sex	TR. Reaction		Cough	Sput.
		Before Med.	After Med.		
W. A.	M	12x15	10x11	-	-
M. E.	M	18x25	14x14	-	-
B. J.	M	18x22	14x15	-	-
F. P.	M	13x14	11x12	-	-
A. G.	M	12x15	8x10	-	-
A. W.	F	65x70	50x60	inc.	inc.
P. M.	F	22x27	22x38	dec.	-
B. J.	F	13x13	21x40	-	-
B. L.	F	22x32	15x40	-	-
C. G.	F	52x60	74x75	-	-

In general, there is no appreciable inhibitory effects after the medication, although four of them showed slight decrease and another four of them showed increase in the size of tuberculin reaction.

Table 2: Short term group

Name	Sex	TR. Reaction		Cough	Sput.
		Before Med.	After Med.		
B. W.	M	12x13	12x12	-	-
L. D.	M	35x42	40x45	-	-
L. S.	M	12x14	12x14	-	-
J. N.	M	18x18	25x35	-	?
L. J.	M	16x16	15x15	-	-
G. M.	F	28x42	45x50	-	-
M. H.	F	32x41	30x40	-	-
P. L.	F	12x14	15x16	-	-
T. J.	F	14x22	20x24	-	-
W. A.	F	18x22	7x11	-	-

In general, there is no appreciable inhibitory effects after the medication, although one of them showed increase in the size of tuberculin reaction.

II. NON-PRE-MEDICATED GROUP

Table 3 shows the results of non-Pre-medicated group.

Table 3: Non-pre-Medicated Group

Name	Sex	TR. Reaction		Cough	Spu
		Before Med.	After Med.		
H. J.	M	23x27	12x12	-	-
C. A.	M	13x13	15x15	-	-
T. J.	M	32x34	28x35	-	-
L. E.	M	28x30	34x45	-	-
S. F.	M	14x22	20x22	-	-
B. A.	F	22x27	20x28	-	-
M. E.	F	14x18	14x16	-	-
R. E.	F	20x24	11x14	-	-
A. N.	F	18x18	10x10	-	-
B. C.	F	17x17	11x12	-	-

There were some inhibitory effect observed that four of ten cases showed considerable decrease and one showed increase in their size of reaction.

III. CONTROL GROUP

Table 4 shows the control group which did not receive any drug but tuberculin test was repeated along with the experimental groups.

Table 4: Control Group

Name	Sex	TR. Reaction		Cough	Spu
		Before Med.	After Med.		
H. J.	M	15x16	13x24	-	-
R. S.	M	18x19	32x42	-	-
D. T.	M	28x55	18x21	-	-
H. F.	M	45x75	45x60	-	-
S. W.	M	26x38	40x70	-	-
T. T.	F	5x7	6x6	-	-
H. L.	F	22x24	18x18	-	-
C. L.	F	15x15	11x11	-	-
N. H.	F	19x19	10x11	-	-
W. P.	F	20x20	55x76	-	-

In general, there is no considerable different in the size of tuberculin reaction in repeated test, although two of them showed decrease and another two of them showed increase in their size of,

skin reaction and remaining of six of them showed essentially unchanged.

DISCUSSION

The advertising campaign to popularize the antihistaminic drugs (10-a, b) has resulted in their extensive use for the treatment of the common cold and general allergic manifestations especially during winter and spring. A correct knowledge regarding the effect of this group of drugs on tuberculin reaction becomes essential. Several articles on this subject both on animal and humans have given rise to contravertial opinions.

As a preliminary study indicated, there has been no appreciable effects on tuberculin reaction one way or the other. However, this study was carried out on the limited small numbers. Therefore, it can be contributory only in a small measure to a final conclusion so that the study was extended to investigate with larger numbers using the most common dosage in practice to minimize the clinical toxic manifestations.

In general, there is no appreciable inhibitory effects on tuberculin reaction considering the variations seen in the size of tuberculin reaction on control group although minor variations observed on each group.

This study left the impression that there is no clinical improvement in the chest condition during or after the medication. Although this conclusion cannot be considered as totally opposed to the findings of Henderson and Judd (3) because they made their observation after giving antihistaminic drug for a period six months and larger doses.

Whereas, in this study, the longest medication was for a 7-day period and the doses were smaller, the author, however, claims that a daily dose, 400 mgm. or more of antihistaminic drugs, was too hazardous because of toxicity.

CONCLUSION

The effects of antihistaminic drugs on the intradermal tuberculin reaction on proved tuberculosis cases was studied.

1. Generally speaking, Pyribenzamine does not play any significant inhibitory effect on tuberculin reaction.
2. Large dose of Pyribenzamine such as daily 400 mgm or more produce undesirable side reactions but promptly disappear following discontinuation of the medication.

Acknowledgement

The author wishes to express his sincere appreciation to Dr. H. S. Willis, Superintendent of the North Carolina Sanatoria for his encouragement and advice.

REFERENCES:-

1. Boquet, A.: Substances antihistaminiques at reactions tuberculiniques, Ann. Inst. Pasteur, 1943, Vol. 69, p. 55.
2. Sarber, R.W.: Effects of Benadryl Hydrochloride on the Tuberculin reaction in Guinea Pigs, Am. Rev. Tuberc., 1948, Vol. 57, P. 504.
3. Judd, A.R. and Henderson, A. R.: The use of antihistaminic drugs in human tuberculosis: A Preliminary Report. Ann. Allergy, 1949, Vol. 7, P. 306.
4. Henderson, A.R.: The Effect of

antihistamine medication on the tuberculin reaction. Am. Rev. Tuberc,

1349, Vol. 60, P. 811.

5. Graub, M. and Barrist, E.: The Effect of Antihistaminic Drugs upon the Tuberculin Reaction. Am. Rev. TB., Vol. 61, P. 735, 1950.
6. Cripe, L.E., Levine, M. L. and Aaron, T. H.: Inhibition of Tuberculin Type of Reaction by Antihistaminic Drugs and Rutin. Am. Rev. TB., 1943, Vol. 52, P. 701.
7. Friedman, E. and Silberman, I.: The Effect of Antihistamine Medication on the Tuberculin Reaction in Children. Am. Rev. TB., 1949, Vol. 60, P. 354.
8. Don Hunter, J. D., Davis and Leroy Hybe: The Effect of Antihistaminic on the Tuberculin Reaction. Am. Rev. TB., 62, 525, 1950.
9. Woobruff, C. E.: Antihistaminics and the Tuberculin Reaction, Am. Rev. TB., 1950, 62, p. 555.
10. a. Editorial: Antihistaminics for Colds, J. A. M. A. Feb. 25, 1950, 142:570.
b. Report of the Council on Pharmacy and Chemistry: Status Report on Antihistaminic Agents in the Prophylaxis and Treatment of Common Cold, J.A.M.A., 142:566-569, Feb. 25, 1950

4-20-51

— 消 息 欄 —

- ① 大韓結核協會 學術大會開催
日時 4287年 11月 中間(日付는追後
公告됨)
場所 慶尙南道馬山市
演題送付處 서울特別市 保健部內

大韓結核協會

本協會 總會 및 學術大會를 上記와如히
開催코지하오나 任員 및 會員 諸位께서는 主
員 參席하여 주심을 바라오며 모-든 苦難
을 무릅쓰고 結核研究에 一踏邁進하고
계시는 諸彦께서도 많은 研究發表가 있
기를 바랍니다

② 結核學會支部設置

今般 本協會에서는 下記場所의 結核學 會
支部를 設置하게 되었음으로 關係諸彦께
서는 많이加入하여 주심을 바랍니다

1. 서울特別市學會支部
2. 釜山學會支部
3. 馬山學會支部
4. 大邱學會支部
5. 光州學會支部

③ 國立中央結核院開院

Model T.B. Center 로서 8月初旬에開
院豫定

部署: 總務部 指導部 治療部 豫防部

場所: 서울特別市 中區會賢洞

會賢洞保健所內

- ④ 서울大學校副總長 尹日善博士은 今般
南興財團에서開催되는 大會議招請으
로 지난7月20日出發

- ⑤ 今般韓美財團으로부터 Full Scholarship
및旅費補助32名을受었다

本協會長 崔昌順氏는 昨年12月 1日

W.H.O. 招請에 依하여 保健社會事業視察
次 歐羅巴等處를 巡廻하고 歸路 巴里 英國
美國等處의 各國 結核協會를 訪問하고
今年 五月 一日에 歸國 하었다

☆	☆	☆
☆		☆
☆		☆
☆	☆	☆