神經症狀로 發病하는 初期結核

國立馬山結核療養所

朴 潤 圭

緒 言

結核療養所를 찾아오는 外來患者中에는 全身倦怠感, 頭痛, 不眠, 眩暈, 心悸亢進, 食慾不振, 不快氣分等 不定한症狀를 呼訴하는 경우가 많았다.

結核發病時는 發熱, 咳嗽, 咯痰, 胸痛等 自覺症狀를 呼訴하는것이 普通이지만 汗便, 不眠, 眩暈, 注意集中不能, 不安感, 記憶障礙等 神經症狀로 結核이發症하는 것도 Pottenger, Pagel, Alexander, Kurt Schlapper, 坂本等 諸氏의 警告한바있다. 이와같은 神經症狀를 呼訴하는患者를 診察한結果 別로 異常을 發見하지못할때는 事實을說明하고, 그대로 물러보내거나, 때로는 神經症으로서 治療하게되는데 이들中에서는 後日에 不幸히도 結核을 發病하는 例를 往往 볼수있다.

主訴	不定한症狀 (全身倦怠, 不眠, 頭痛等)
檢 査 成 績	Tuberculin反應(一)~(十) 血沈 正常 X線 檢 査 理學的檢査

↓
神經衰弱으로 診斷

↓
放置 또는 加療

↓
結核發病

이와같은症狀를 呼訴하는 患者를 單純히 神經症으로만 生覺한수없으며, 肺結核의 前驅症狀이 아닌가도 疑心해볼 必要가 있다.

東京慈大 古閑氏는 이와같은 狀態를 별써부터 注目하고 結核을 原因으로하는 이와같은 症狀群을 Camouflaged Tuberculosis with Neurotic Syndrome(迷彩結核)이라고 부르고

있는데 單純한 神經症과 區別하여 結核을 早期에 發見하는데 便宜한 檢査方法을 發表하고 있다.

鑑別할 疾患

上述한 神經症狀는 急性 또는 慢性疾患, 或은 神經症이나 精神病의 初期에 共通인 것으로 그呼訴하는 症狀만으로 判斷하는것은 매우危險하며 慎重히 檢討하여 正確한診斷이 決定되어야 할것이다. 鑑別할疾患으로서 高血壓, 甲狀腺機能亢進, 脚氣, 寄生虫症, 糖尿病, 腎臟炎, 輕症敗血症, 梅毒, 貧血, 惡性腫瘍, 結核, 神經症, 低血壓, 藥物中毒, Vitamin缺乏症等 이있다.

臨床의 諸像

上記 古閑敎室에서 約 2年동안에 取扱한 77例(男50, 女27)에對한 觀察한結果를 紹介하면 다음과같다.

1) 年齡

男子는 14歲 乃至 38歲 平均 22歲

女子는 16歲 乃至 41歲 平均 27歲

結核發動年齡에 該當하는것이 興味깊은 일이다.

2) 主訴

全身倦怠感	21(27.1%)
頭重	15(19.4%)
不眠	14(18.1%)
食慾不振	14(18.1%)
頭痛	11(14.2%)
眩暈	11(14.2%)
心窩部不快感	10(12.9%)
心悸亢進	10(12.9%)
微熱	8(10.3%)

3) 튜버큐린反應

結核感染의 臨床的證明은 튜버큐린反應

이 第一 混淆하다는것은 周知의事實이지만, 이點에關해서는 아직 疑問이 多少있다. 重症 結核에서 가끔 陰性을 나타내는것은 널리 알려진 事實이고, 近藤氏는 50歲以上の 老人結核에있어 半數가 陰性임을 報告하고있으며, 大澤 清水 兩氏는 開放性結核 約 25%에서 陰轉을 觀察하고있다. 上記77名의 成績은

陽性者(陽轉期判明)	48名
(陽轉期不明)	10名
疑陽性者	6名
陰性者	13名

以上과같으나, 陰性이라도 注意해야된다.

4) 血沈

77名中 11例의輕度促進을 除外하고는 全部 正常이었는데 우리들의 日常臨床에서도 X線上 廣範한陰影이 認定될때라도 靜止狀態에 있을때는 血沈이 促進하지 않는것을 자주 볼수있다.

(參考) 肺結核各病型의血沈値(Thiele)

病型	血沈値 正常(1~5mm)	高度促進 (30mm以上)
硬化型	66.8%	2.1%
硬化增殖型	45.6%	7.0%
硬化空洞型	24.3%	13.1%
主로增殖型	15.5%	21.5%
滲出型	5.4%	49.1%

5) 結核菌

少量의痰을 自覺하는者가 17例었는데 塗沫, 培養 모두 陰性이였으며, 이 17例를 包含한23例에서 早朝空腹時의 胃液培養結果도 모두 陰性이었다.

이 Camouflaged Tuberculosis의狀態는 菌排出을 하지않는程度의 輕度の病變을 想定할수있지않을가 生覺된다.

6) 微熱

微熱을 呼訴한例는 8例인데 初診時 微熱이 發見된者 35名(45.5%)으로서 매우 重要한症狀으로 生覺된다.

7) X線所見

從來의觀點에서는 當然히 正常範圍內에있다고 生覺되는데 詳細히 吟味해본結果는 다

음과 같다. (65例)

1. 肺門腺腫大의印象을주는者	13
2. 橫隔膜肋膜炎의痕跡있는者	10
3. 肺門陰影增強	9
4. 肺紋理增強	6
5. 毛髮像(Haerlinie)	3
6. 縱隔竇肋膜炎의 所見있는者	7
7. 石灰沈着	3
8. 初感染巢	1

全然 疑問을 품을수없는正常者는 11(16.9%)例였다.

Gvaeft, Kupferle等은 一定한크기 以下의病巢는 X線檢査로서도 證明할수없으며, 완두大의病巢도 그位置에따라서는 잘나타나지 않는다고 말하고있다.

8) 向性調査

古閑氏向性調査表에依함

疾患名	內向性	外向性	中間型	計
迷彩結核	37(67.3%)	11(20.0%)	7(12.7%)	55
肺結核(鈴木)	172(61.4%)	75(26.8%)	33(11.8%)	280
神經症(古閑)	36(90.0%)	4(10.0%)	0	40
Pottenger	10(58.8%)	5(29.4%)	2(11.7%)	17
馬山結核療養所	128(65.6%)	41(21.0%)	26(13.3%)	195

9) 機能病理

Pottenger氏는 結核症狀을 3群으로 分類하고있는데 第一群은 中毒症狀
第二群은 內臟反射症狀
第三群은 病巢局所症狀이다.

區別	疾患	迷彩結核	肺結核(鈴木)	馬山療養所
第一群		78.4%	27.4%	41.9%
第二群		14.9%	33.5%	46.7%
第三群		6.0%	36.5%	47.7%

肺結核과는 反對의關係를 나타내고있다.

10) 下垂體副腎系機能

(流血中好酸球數의比較)

正常値70~450/mm³ (Kark, Best)

健康人 225 · 肺結核 372

迷彩結核 373 · 神經症 62

診 斷

아직도 決定的檢査方法은 없으나 上述한 여러가지病像은 參考로하는데 下記의7項目이 目標가된다.

1. 精神傾向의洞察
2. 呼訴하는症狀이單純한것
3. 다른 神經症 症狀을 많이 合併치않은것
4. 류버큐린反應의 吟味 (陰性이라도注意)
5. 微熱
6. X線所見은 從來 結核을 否定하고 正常範圍로 生覺되는데도 慎重히 檢討할 것
7. 流血中好酸球數

發 現 機 轉

이와같은精神神經症狀의發現은 主로 初感染 時에 나타난다고 生覺되는데, 結核菌毒素의 植物神經系統 或은 어떤內分泌系統에의 親和性에 因한다고 說明되고있으나 이것이 모든患者

에서 出現하는것도 아니고 그事態는 아직 不明한 點이 많다.

이와같이 精神神經症狀이 著明하게 나타나는 境遇는 特殊性格 即 神經質素質에 起因한다고 生覺하는것이 便利한듯하다. 이性格의特徵으로서 內向性和 完全慾이 強하기때문에 普通人이 看過해버릴 程度의 輕微한症狀(初期結核의中毒症狀)에도 注意가 固着하여 不安感을 일으키고 漸漸 病覺을 擴大伸張시켜 當初에 想像도 못했던 病像을 만들었다고 生覺된다.

治 療

安靜, 榮養을 主로하는 療養指導을하고 必要에 따라서는 化學療法, 對症療法을 한다.

77例中 安靜榮養만으로	32例
INAH投與	28例
비타민B投與	7例
對症療法	8例
SM, PAS	2例

以上

Early Tuberculosis with Neurotic Symptoms

YUN KYU, PAK M. D.

Masan National T.B. Sanatorium.

"Introduction"

Among the out-patients who visit a TB sanatorium, we find some patients who complain of indefinite symptoms such as headache, insomnia, dizziness, palpitations, loss of appetite, and malaise.

At the onset of tuberculosis, patients usually complain of fever, cough, sputum and chest pain, but on the other it was mentioned by Pottenger, Pagel, Alexander, Kurs-Schlapper and Sakamoto that tuberculosis often begins with neurotic symptoms such as insomnia, dizziness, anxiety, disturbance of memory and dis-

turbance of attention and concentration.

After examining those out-patients who complain of neurotic symptoms, and finding no organic changes, we usually explain to them that there is nothing wrong with them.

Sometimes we try to treat them for neurosis. But we recognize that some of those out-patients unfortunately develop tuberculosis in the future. It is unwise to make a diagnosis of neurosis in such out-patients, and possibility of their symptoms being those of early tuberculosis must always be born in mind.

Dr Koga, Professor of Tokyo Jikeikai School of Medicine, has called attention previously to such symptoms which may be tuberculous in origin and refers to such a condition as "Camouflaged tuberculosis with neurotic syndrome" as previously reported together with methods of examination which helps to differentiate the two conditions of neurosis and early tuberculosis.

Seventy-seven so-called cases of camouflaged tuberculosis patients were investigated at Koga Clinic for 2 years (Aug 1952~June 1954)

„Clinical Picture"

(1) Age

Male 14 to 38 Averagd 22

Female 16 to 41 Average 27

It may be significant that the ages of those patients are similar to those of tuberculosis onset.

(2) Chief complaint.

General malaise	21(27.1%)
Heavy sensation of head	15(19.4%)
Insomnia	14(18.1%)
Loss of appetite	14(18.1%)
Headache	11(14.1%)
Dizziness	11(14.1%)
Uncomfortable sensation	
in the epigastrium	10(11.9%)
Palpitations	10(11.9%)
Fever	8(10.3%)

(3) Tuberculin Reaction

It is a well known fact that tuberculin testing is the best way clinically to prove the tuberculous infection, but we still have an unsolved problem concerning Tuberculin reaction.

Among the above 77 patients, 13 negative and 6 false positive were reported.

(4) Sed. rate

Exept for a slight increase in 11 pa

tients, the rest were all normal.

(5) AFB

17 patients complained of sputum and examination, both smear and culture for AFB on those patients were negative.

13 cultures(including the above 17) of gastric washings all showed negative results.

(6) Slight fever

8 Patients complained of fever. It was found that at the first interview 35 had slight fever, so this fever is significant in making diagnosis.

(7) X-Ray findings

They are all within normal limit, but the following findings are seen on close observation.

- Suspicious enlargement of hilar gland
- Evidence of phrenicopleuritis
- Increased hilar shadow
- Increased lung markings
- Haarlinie
- Increased opacity lateral to Mediastinal shadow
- Calcification

(8) Eosinophile Count

Normal.....225

Camou flagged TB.....275

TB.....375

Neurosis.....62

Normal 70 to 450/mm3
(Kark, Best)

"Pathogenic Mechanism"

These indefinite symptoms seem to appear mainly in the early stages of tuberculous infection. It is sometimes given as an explanation that the causative agent is the toxin of tubercle bacilli, but the exact pathogenic mechanism has not been clearly explained. It may be convenient to explain by differences in personality. A nervous person easily feels slight

psychosomatic changes which most people do not notice. The perpetuation of these symptoms due to nervousness must also be emphasized.

‘Treatment’

The treatment camouflaged tuberculosis is the same as that of overt tuberculosis, bed rest, good nutrition, symptomatic treatment and chemotherapy if necessary.

Among the 77 patients;

32 treated with bedrest and good nutrition

28 treated with INH

7 treated with Vitamin B

8 treated symptomatic

2 treated with SM and PAS.

All these patients responded to this treatment, and became symptom free. More developed overt tuberculosis.